Jan. 3, 1984

[54] METHOD FOR THE REDUCTION OF MUCIN VISCOSITY

[75] Inventors: Anthony Cerami, Flanders, N.J.; Nina F. Tabachnik, Little Neck, N.Y.

[73] Assignee: **The Rockefeller University, New** York, N.Y.

[21] Appl. No.: 62,503

[22] Filed: Jul. 31, 1979

[51] Int. Cl.³ A01N 31/66; A01N 31/15; A01N 31/44; A01N 31/47

[56] References Cited

U.S. PATENT DOCUMENTS

3,892,824 7/1975 Piper et al. 424/211

OTHER PUBLICATIONS

"Studies on the Reduction of Sputum Viscosity in Cystic Fibrosis Using an Orally Absorbed Protected

Thiol", Tabachnik, Peterson, and Cerami, Apr. 22, 1980; vol. 214, No. 2; pp. 246-249.

"Protein Binding of N-2-Mercaptoethyl-1,3-Diaminopropane via Mixed Disulfide Formation after Oral Administration of WR 2721¹", Tabachnik, Blackburn, Peterson and Cerami; vol. 220, No. 2; Oct. 19, 1981; pp. 243-246.

"Biochemical and Rheological Characterization of Sputum Mucins from a Patient with Cystic Fibrosis", Tabachnik, Blackburn, and Cerami; vol. 256, No. 14; Jul. 25, 1981, pp. 7161–7165.

Survey of Compounds which have been Tested for Carcinogenic Activity, pp. 21,23,113,153,154,469; Thompson and Co., Publishers, DHEW Pub. No. (NIH)72-35 (1968-1969 volume).

Primary Examiner—Alan L. Rotman Attorney, Agent, or Firm—Oblon, Fisher, Spivak, McClelland & Maier

[57] ABSTRACT

A method for reducing mucin viscosity which comprises administering an effective dose of a compound having protected sulfhydryl groups which metabolize in vivo to produce free sulfhydryl groups.

8 Claims, 3 Drawing Figures



